

$B_{s,d} \rightarrow \gamma\gamma$ Decay with the Fourth GenerationWujun Huo^a, Cai-Dian Lü^b, and Zhenjun Xiao^c*a. Department of Physics, Peking University, Beijing 100871, P.R. China**b. CCAST (World Laboratory), P.O.Box 8730, Beijing 100080, P.R.China;**b. Institute of High Energy Physics, CAS, P.O. Box 918(4), Beijing 100039, P.R.China***c. Physics Department, Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China*

(February 1, 2008)

Abstract

We calculate the branching ratios of $B_{s,d} \rightarrow \gamma\gamma$ decay in the sequential fourth generation model. We find that the theoretical values of the branching ratios, $\text{BR}(B_s \rightarrow \gamma\gamma)$, including the contributions of t' , are much different from the minimal standard model (SM) predictions. The new physics contribution can provide one to two orders enhancement to the standard model prediction of $\text{BR}(B_s \rightarrow \gamma\gamma)$. But due to the tiny values of 4th generation CKM matrix element $V_{t'd}^* V_{t'b}$, the new physics effect on $\text{BR}(B_d \rightarrow \gamma\gamma)$ is very small and can not be distinct from the SM prediction. It is shown that the decay $B_s \rightarrow \gamma\gamma$ can give the test of new physics signals from the 4th generation model.

Typeset using REVTeX

*Mailing address

I. INTRODUCTION

The minimal Standard Model (SM) is a very successful theory of the elementary particles known today. But it should not be the final theory, because it has too many unknown parameters to be put by hand. Most of these parameters are in the fermion part of the theory. We don't know the origin of the quarks and leptons, as well as how to determinate their mass and number theoretically. We have to get their information all from experiments. There is still not a successful theory which can unify all the four basic interactions. Perhaps our *elementary particles* have substructure and we need a more fundamental theory than SM. This is beyond the scope of our current experiments. On the other hand, the recent measurement of the muon anomalous magnetic moment by the experiment E821 [1] disagrees with the SM expectations at more than 1.6σ - 3σ [2,3] level. Moreover, there are convincing evidences that neutrinos are massive and oscillate in flavor [4]. It seems to indicate the presence of new physics. The discovery of neutrino oscillation has been one of the most exciting experimental results in the recent years [5]. As the experimental data accumulating in the Super-Kamiokande collaboration [6,7], more results published by SNO [8], K2K [9] and CHOOZ [10] experiments. These experiments give possible signals of new physics beyond SM.

In this paper, we consider the sequential fourth generation model to estimate the possible contributions to the decay $B_{s,d} \rightarrow \gamma\gamma$ from the exchange of the fourth up-type quark t' . From the point of phenomenology, there is a realistic question one may ask: What is the number of the fermion generations or whether there are additional quarks or leptons other than the known three generations. The present experiments can tell us there are only three generation of fermions with *light* neutrinos whose mass is smaller than $M_Z/2$ [11], but the experiments don't exclude the existence of other kind of additional generation, such as the fourth generation, with a *heavy* neutrino, i.e. $m_{\nu_4} \geq M_Z/2$ [12]. Latest studies in the electroweak sector allow the existence of a fourth generation with heavy Dirac neutrino [13]. Many authors studied models which extend the fermions part, such as vector-like quark models [14], sterile neutrino models [15] and the sequential four generation standard model (SM4) [16]. Here we consider the sequential fourth generation, non -supersymmetric model, which includes an up-like quark t' , a down-like quark b' , a lepton τ' , and a heavy neutrino ν' besides the known three generation of fermions in the SM. The properties of these new fermions are the same as their corresponding counterparts of other three generations except their masses and CKM matrix elements.

Many attempts have been made recently about the fourth generation [17–19] and the experimental searches of the fourth generation particles [20,21]. In our previous papers [22–24], we investigated the rare B meson decays with the fourth generation [22], ϵ'/ϵ in K^0 systems and $\Delta M_{B_{s,d}}$ in SM4 [23]. We got some interesting results, such as the new effects of the 4th generation particle on the meson decays and CP violation. We also got the constraints of the fourth generation CKM matrix factors [22,23]. Recently, these decays in fourth generation were considered again. In ref. [17], the phases of the fourth generation CKM matrix factors, $V_{t's}^* V_{t'b'}$, have been included. The exclusive B decays with the fourth generation were investigated in Ref. [18]. These rare decays are very interesting to study the new physics effects of the fourth generation. In this paper, we would like to investigate the rare decay $B_{s,d} \rightarrow \gamma\gamma$ with the fourth generation.

As a loop-induced flavor changing neutral current (FCNC) process the inclusive decay (at quark level) $b \rightarrow s\gamma\gamma$ is in particular sensitive to contributions from those new physics beyond the SM. There is a vast interest in this decay. On the experimental side, only upper limits (90%*C.L.*) on the branching ratios of $B_{s,d} \rightarrow \gamma\gamma$ decay are currently available

$$\text{BR}(B_s \rightarrow \gamma\gamma) < 1.48 \times 10^{-4}, \quad [25], \quad (1)$$

$$\text{BR}(B_d \rightarrow \gamma\gamma) < 1.7 \times 10^{-6}, \quad [26], \quad (2)$$

which are roughly two orders above the SM predictions [27–29],

$$5 \times 10^{-7} < \text{BR}(B_s \rightarrow \gamma\gamma) < 15 \times 10^{-7}, \quad (3)$$

$$1 \times 10^{-8} < \text{BR}(B_d \rightarrow \gamma\gamma) < 4.0 \times 10^{-8}. \quad (4)$$

Within the SM, the electroweak contributions to $b \rightarrow s\gamma\gamma$ and $B \rightarrow \gamma\gamma$ decays have been calculated some time ago [27], the leading-order QCD corrections and the long-distance contributions were evaluated recently by several groups [28,30]. The new physics corrections were also considered, for example, in the two-Higgs doublet model [31,32] and the supersymmetric model [33]. There is also study concerning other exclusive channels [34]. In a previous paper [35], we calculated the Technicolor corrections to the rare decays $B_{s,d} \rightarrow \gamma\gamma$ in the QCD factorization approach. The great progress in theoretical studies recently encourage us to do more investigations about this decay in the sequential fourth generation model.

This paper is organized as the following: In Section 2, we present the formulae for the decays $B_{s,d} \rightarrow \gamma\gamma$ in SM and the fourth generation model. We also obtain the constraints of the fourth generation CKM matrix factors, $V_{t'd}^*V_{t'b}$ and $V_{t's}^*V_{t'b}$ from experiments in this section. In Section 3, we give the numerical analysis of the branching ratios in the fourth generation model. The conclusions are in the last section.

II. $B_{S,D} \rightarrow \gamma\gamma$ IN THE FOURTH GENERATION MODEL

Same as in our last paper [35], up to the corrections of order $1/m_W^2$, the effective Hamiltonian for $b \rightarrow s\gamma\gamma$ at scales $\mu_b = \mathcal{O}(m_b)$ is just the one for $b \rightarrow s\gamma$ and takes the form

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \left[\sum_{i=1}^6 C_i(\mu_b) Q_i + C_{7\gamma}(\mu_b) Q_{7\gamma} + C_{8G}(\mu_b) Q_{8G} \right], \quad (5)$$

where in view of $|V_{us}^* V_{ub}/V_{ts}^* V_{tb}| < 0.02$, we have neglected the term proportional to $V_{us}^* V_{ub}$. Here $Q_1 \dots Q_6$ are the usual four-fermion operators whose explicit form is given below. The last two operators in the Eq.(5), characteristic for this decay, are the *magnetic-penguin* operators. The complete list of operators is given as follows

$$O_1 = (\bar{c}_{L\beta} \gamma^\mu b_{L\alpha}) (\bar{s}_{L\alpha} \gamma_\mu c_{L\beta}), \quad (6)$$

$$O_2 = (\bar{c}_{L\alpha} \gamma^\mu b_{L\alpha}) (\bar{s}_{L\beta} \gamma_\mu c_{L\beta}), \quad (7)$$

$$O_3 = (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta} \gamma_\mu q_{L\beta}), \quad (8)$$

$$O_4 = (\bar{s}_{L\alpha}\gamma^\mu b_{L\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta}\gamma_\mu q_{L\alpha}) , \quad (9)$$

$$O_5 = (\bar{s}_{L\alpha}\gamma^\mu b_{L\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta}\gamma_\mu q_{R\beta}) , \quad (10)$$

$$O_6 = (\bar{s}_{L\alpha}\gamma^\mu b_{L\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta}\gamma_\mu q_{R\alpha}) , \quad (11)$$

$$O_7 = (e/16\pi^2)m_b\bar{s}_L\sigma^{\mu\nu}b_RF_{\mu\nu} , \quad (12)$$

$$O_8 = (g/16\pi^2)m_b\bar{s}_L\sigma^{\mu\nu}T^ab_RG_{\mu\nu}^a . \quad (13)$$

It is the magnetic γ -penguin operator O_7 , which plays the crucial role in this decay. The effective Hamiltonian for $b \rightarrow d\gamma\gamma$ is obtained from Eqs.(5-13) by the replacement $s \rightarrow d$.

In the case of the fourth generation there is an additional contribution to $b \rightarrow (s, d)\gamma\gamma$ from the virtual exchange of the fourth generation up quark t' . In Fig.1, we draw the relevant Feynman diagrams which contribute to the decays $b \rightarrow (s, d)\gamma\gamma$. The Wilson coefficients of the dipole operators are given by

$$C_{7,8}^{\text{eff}}(\mu_b) = C_{7,8}^{(\text{SM})\text{eff}}(\mu_b) + \frac{V_{t'q}^* V_{t'b}}{V_{tq}^* V_{tb}} C_{7,8}^{(4)\text{eff}}(\mu_b), \quad (14)$$

where $C_{7,8}^{(4)\text{eff}}(\mu_b)$ represent the contributions of t' to the Wilson coefficients. We recall here that the CKM coefficient corresponding to the t quark contribution, i.e., $V_{ts}^* V_{tb}$, is factorized in the effective Hamiltonian as given in Eq. (5). The formulae for calculating the Wilson coefficients $C_{7,8}^{(4)}(m_W)$ are the same as in the SM except exchanging t' quark for t quark.

In SM4, the quark mixing matrix can be written as,

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} & V_{ub'} \\ V_{cd} & V_{cs} & V_{cb} & V_{cb'} \\ V_{td} & V_{ts} & V_{tb} & V_{tb'} \\ V_{t'd} & V_{t's} & V_{t'b} & V_{t'b'} \end{pmatrix}, \quad (15)$$

where $V_{qb'}$ and $V_{t'q}$ are the elements of the 4×4 CKM mixing matrix of the SM4, which now contains nine parameters, i.e., six angles and three phases. The rest elements without prime in the matrix are the usual three generation CKM matrix elements.

The current experimental bounds of $V_{t's}^* V_{t'b}$ and $V_{t'd}^* V_{t'b}$ can be found in ref. [24]. The factor $V_{t's}^* V_{t'b}$ can be constrained by the decay $B \rightarrow X_s \gamma$ [22,24]

$$-11.6 \times 10^{-2} < V_{t's}^* V_{t'b}^{(1)} < -6.1 \times 10^{-2}, \quad (16)$$

or

$$1.9 \times 10^{-3} < V_{t's}^* V_{t'b}^{(2)} < 3.6 \times 10^{-3}. \quad (17)$$

Considering the unitarity conditions of the CKM matrix,

$$\sum_i V_{is}^* V_{ib} = 0, \quad (i = u, c, t, t'), \quad (18)$$

and taking the average values of the SM 3×3 CKM matrix elements from Ref. [36], we obtain,

$$|V_{t's}^* V_{t'b}| < 7.6 \times 10^{-2}. \quad (19)$$

Combining the above constraint, we arrive at

$$-7.6 \times 10^{-2} < V_{t's}^* V_{t'b}^{(1)} < -6.0 \times 10^{-2}, \quad (20)$$

$$1.9 \times 10^{-3} < V_{t's}^* V_{t'b}^{(2)} < 3.6 \times 10^{-3}. \quad (21)$$

In the following numerical calculation, we take the values $V_{t's}^* V_{t'b}^{(1)} = -7.0 \times 10^{-2}$ and $V_{t's}^* V_{t'b}^{(2)} = 2.5 \times 10^{-3}$, for illustration.

For the CKM factor $V_{t'd}^* V_{t'b}$, we can get its constraint from the present experimental value of ΔM_{B_d} [24]. Also considering the unitarity conditions of the CKM matrix,

$$\sum_i V_{id}^* V_{ib} = 0, \quad (i = u, c, t, t'), \quad (22)$$

we finally get

$$-1.0 \times 10^{-4} < V_{t'd}^* V_{t'b} < 0.5 \times 10^{-4}. \quad (23)$$

In the next section, we will take the values $V_{t'd}^* V_{t'b} = -1.0 \times 10^{-4}$, -0.5×10^{-4} , -0.1×10^{-4} , 0.1×10^{-4} and 0.5×10^{-4} in numerical calculation respectively.

III. THE $B_{S,D} \rightarrow \gamma\gamma$ DECAY RATES AND PHENOMENOLOGY

We first calculate the decay $B_s \rightarrow \gamma\gamma$ (the formulas in the case of $B_d \rightarrow \gamma\gamma$ can be obtained just by the replacement of $s \rightarrow d$). For this exclusive decay of B_s meson, we need to deal with both of the long distance and short distance QCD corrections in the B_s meson side. Applying the factorization assumption, we shall adapt a phenomenological approach where the long distance effects are replaced by a few non-perturbative parameters. In other words we simply evaluate the hadronic matrix element of $M_I + M_R$ from one particle irreducible (1PI) and one particle reducible (1PR) Feynman diagrams, relying on a phenomenological model. Before doing so, it is important to note that M_R is apparently non-local due to internal b or s quark propagator. To handle these non-local terms, one observes that the b quark inside the B_s meson carries most of the meson energy, and its four velocity can be treated as equal to that of B_s . Hence one may write b quark momentum as $p = m_b v$ where v is the common four velocity of b and B_s . With this parametrization, we have

$$\begin{aligned} p \cdot k_1 &= m_b v \cdot k_1 = \frac{1}{2} m_b m_{B_s} = p \cdot k_2, \\ p' \cdot k_1 &= (p - k_1 - k_2) \cdot k_1 = -\frac{1}{2} m_{B_s} (m_{B_s} - m_b) = p' \cdot k_2, \end{aligned} \quad (24)$$

where the second equation is based on a constituent picture [27] that b and $\bar{s}$ quarks share the total energy of B_s ¹. Therefore $-p'$ should be taken as the four momentum of the constituent

¹Note that the momentum of $\bar{s}$ quark is $-p'$ as p' denotes the momentum of s quark in $b \rightarrow s\gamma\gamma$.

$\bar{s}$ quark. With Eq.(24), M_R is readily made local. We then compute the amplitude for $B_s \rightarrow \gamma\gamma$ using the following relations

$$\begin{aligned}\langle 0|\bar{s}\gamma_\mu\gamma_5 b|B_s(P)\rangle &= -if_{B_s}P_\mu, \\ \langle 0|\bar{s}\gamma_5 b|B_s(P)\rangle &= if_{B_s}M_B,\end{aligned}\tag{25}$$

where f_{B_s} is the B_s meson decay constant which is about 200 MeV according to recent Lattice QCD calculations [37].

The total amplitude is now separated into a CP-even and a CP-odd part

$$T(B_s \rightarrow \gamma\gamma) = M^+ F_{\mu\nu} F^{\mu\nu} + iM^- F_{\mu\nu} \tilde{F}^{\mu\nu}.\tag{26}$$

We find that

$$M^+ = -\frac{4\sqrt{2}\alpha G_F}{9\pi} f_{B_s} V_{ts}^* V_{tb} \left(\frac{m_b}{m_{B_s}} B K(m_b^2) + \frac{3C_7}{8\bar{\Lambda}} \right),\tag{27}$$

with $B = -(3C_6 + C_5)/4$, $\bar{\Lambda} = 1 - m_b/m_{B_s}$, and

$$M^- = \frac{4\sqrt{2}\alpha G_F}{9\pi} f_{B_s} V_{ts}^* V_{tb} \left(\sum_q A_q J(m_q^2) + \frac{m_b}{m_{B_s}} B L(m_b^2) + \frac{3C_7}{8\bar{\Lambda}} \right),\tag{28}$$

where

$$\begin{aligned}A_u &= (C_3 - C_5)N_c + (C_4 - C_6), \\ A_d &= \frac{1}{4} [(C_3 - C_5)N_c + (C_4 - C_6)], \\ A_c &= (C_1 + C_3 - C_5)N_c + (C_2 + C_4 - C_6), \\ A_s &= A_b = \frac{1}{4} [(C_3 + C_4 - C_5)N_c + (C_3 + C_4 - C_6)].\end{aligned}\tag{29}$$

The functions $J(m^2)$, $K(m^2)$ and $L(m^2)$ are defined by

$$\begin{aligned}J(m^2) &= -\frac{1}{2} + \frac{1}{z} I_{00}(m^2), \\ K(m^2) &= -2 + \frac{4-z}{z} I_{00}(m^2), \\ L(m^2) &= I_{00}(m^2),\end{aligned}\tag{30}$$

where $z = m_{B_s}^2/m^2$, and

$$I_{00}(m^2) = -\int_0^1 dx \frac{1}{x} \ln(1 - zx + zx^2)\tag{31}$$

$$= \begin{cases} 2 \arctan^2(\sqrt{\frac{z}{4-z}}); & z < 4 \\ \frac{\pi^2}{2} - 2 \log^2(\frac{\sqrt{z} + \sqrt{z-4}}{2}) + 2i\pi \log(\frac{\sqrt{z} + \sqrt{z-4}}{2}); & z > 4 \end{cases}.\tag{32}$$

The decay width for $B_s \rightarrow \gamma\gamma$ is simply

$$\Gamma(B_s \rightarrow \gamma\gamma) = \frac{m_{B_s}^3}{16\pi}(|M^+|^2 + |M^-|^2). \quad (33)$$

To obtain numerical results, we have set light quark masses to zero and used [36] $m_t = 175$ GeV, $m_b = 5.0$ GeV and $m_c = 1.5$ GeV. Furthermore, we take $m_{B_s} = 5.37$ GeV, $m_{B_d} = 5.28$ GeV and $\alpha = 1/129$. The numerical values for Wilson coefficients $C_1 - C_8$ evaluated in SM at $\mu = m_b$ are listed in Table I.

We first consider the decay mode $B_s \rightarrow \gamma\gamma$. Figs.2, 3 and 4 show the branching ratio $Br(B_s \rightarrow \gamma\gamma)$ *versus* the mass of t' and the fourth generation CKM matrix factors $V_{t's}^* V_{t'b}^{(1)}$ and $V_{t's}^* V_{t'b}^{(2)}$, respectively. From these figures, we find that the SM4 predictions for the branching ratio $Br(B_s \rightarrow \gamma\gamma)$ are rather different from that of the standard model. The decay rates can be enhanced by about 1-2 orders relative to the SM prediction in the reasonable range of $m_{t'}$ and $V_{t's}^* V_{t'b}$. This implies that the new physics signal of the fourth generation Model may come from the decay of $B_s \rightarrow \gamma\gamma$. The future experiment of LHCb in CERN can test it. From Fig.2, we find that the branching ratio of $B_s \rightarrow \gamma\gamma$ increase with the mass of t' for the fourth generation CKM matrix factor, $V_{t's}^* V_{t'b} = -7.0 \times 10^{-2}$ and 2.5×10^{-3} . When $V_{t's}^* V_{t'b} = -7.0 \times 10^{-2}$, the curve increase rapidly with $m_{t'}$. But when $V_{t's}^* V_{t'b} = 2.5 \times 10^{-3}$, the branching ratio changes very slowly. Because the fourth generation CKM matrix factor, $V_{t's}^* V_{t'b}$, is small in the latter case. It can be seen from Eq. (14) that the new physics contribution is proportional to the size of the CKM factor $V_{t's}^* V_{t'b}$, thus it is very important in the new physics contribution with $m_{t'}$. From Figs.3 and 4, one can also see that, the branching ratio of $B_s \rightarrow \gamma\gamma$ decrease with $V_{t's}^* V_{t'b}^{(1)}$ but increase with $V_{t's}^* V_{t'b}^{(2)}$, since $V_{t's}^* V_{t'b}^{(1)}$ is negative and gives the opposite contributions as SM to the coefficients $C_{7,8}^{eff}$ (see Eq. (14)).

For the decay $B_d \rightarrow \gamma\gamma$, it is easy to find from Figs.5 and 6 that the new physics contribution to the branching ratio is tiny. It is shown that it is not suitable to test the new physics signals from the 4th generation in this channel. The prediction of the branching ratio in the 4th generation model is between 2.35×10^{-8} and 2.45×10^{-8} , when we take the reasonable values of $m_{t'}$ (from 200 GeV to 1000 GeV) and $V_{t'd}^* V_{t'b}$ (see Eq. (23)). This numerical result is fully in the range of the SM prediction (see Eq. (4)). The reason is that the experimental constraint of the 4th generation CKM matrix element factor $V_{t'd}^* V_{t'b}$ is very stringent (only at 10^{-4} level, see Eq.(23)). We also find that the curves change little along with the variations of $m_{t'}$ and $V_{t'd}^* V_{t'b}$. The reason is also from the tiny $V_{t'd}^* V_{t'b}$. We can see that the branching ratios of both $B_s \rightarrow \gamma\gamma$ and $B_d \rightarrow \gamma\gamma$ decays strongly depend on the size of the related mixing elements.

IV. CONCLUSION

As a summary, the size of new physics contributions to the rare decays of $B_{s,d} \rightarrow \gamma\gamma$ from the fourth generation strongly depends on the values of the masses of the new heavy up quark t' and the 4th generation CKM matrix elements, $V_{t's}^* V_{t'b}$ and $V_{t'd}^* V_{t'b}$. The results in the 4th generation model are quite different from the SM case for decay $B_s \rightarrow \gamma\gamma$, its branching ratio can be enhanced by about 1-2 orders of magnitude. A large branching ratio of $B_s \rightarrow \gamma\gamma$ decay measured in experiment may be a new physics signal of the 4th generation model. But for decay $B_d \rightarrow \gamma\gamma$, because of the tiny values of 4th generation CKM matrix

element, $V_{t'd}^* V_{t'b}$, the new physics contribution to $\text{BR}(B_d \rightarrow \gamma\gamma)$ is very small and can not be distinct from the SM prediction.

ACKNOWLEDGMENT

W.J. Huo acknowledges the support from the Chinese Postdoctoral Science Foundation. from the Natural Sciences and Engineering Research Council of Canada. C.D. Lü acknowledges the support by National Science Foundation of China under Grants No. 90103013 and 10135060. And Z.J. Xiao acknowledges the support by the National Natural Science Foundation of China under Grants No. 10075013 and 10275035, and by the Research Foundation of Nanjing Normal University under Grant No. 214080A916.

REFERENCES

- [1] Muon g-2 Collaboration, H.N. Brown *et al.*, Phys. Rev. Lett. **86**, 2227 (2001).
- [2] U. Chattopadhyay, A. Corille and P. Nath, hep-ph/0202275; M. Knecht and A. Nyffeler, hep-ph/0111058; M. Knecht, A. Nyffeler, M. Perrottet and E. de Pafael, Phys. Rev. Lett. **88**, 071802 (2002); M. Hayakawa and T. Kinsoshita, hep-ph/0112102; I. Blokland, A. Czarnecki and K. Melnikov, Phys. Rev Lett. **88**, 071803 (2002); J. Bijnens, E. Pallante and J. Prades, hep-ph/0112255.
- [3] K. Hagiwara, A. D. Martin, Daisuke Nomura and T. Teubner, hep-ph/0209187.
- [4] Y. Fukuda *et al.*, Phys. Lett. **B436** 33 (1998); Phys. Rev. Lett. **81**, 1562 (1998).
- [5] Y. Fukuda *et al.*, Super-Kamiokande Collaboration, Phys. Rev. Lett. **81**, 1562 (1998); **82**, 2644 (1999); **82**, 1810 (1999); **82**, 2430 (1999).
- [6] Y. Fukuda *et al.*, Super-Kamiokande Collaboration, Y. Fukuda *et al.*, Super-Kamiokande Collaboration, Phys. Rev. Lett. **86**, 5651 (2001).
- [7] Y. Fukuda *et al.*, Super-Kamiokande Collaboration, Phys. Rev. Lett. **86**, 5656.
- [8] Q. R. Ahmad *et al.*, SNO Collaboration, Phys. Rev. Lett. **87**, 071301 (2001).
- [9] S. H. Ahn *et al.*, K2K Collaboration, Phys. Lett. **B511**, 178 (2001).
- [10] M. Apollonio *et al.*, CHOOZ Collaboration, Phys. Lett. **B420**, 397 (1998); **B466**, 415 (1999).
- [11] G.S. Abrams *et al.*, Mark II Collab., Phys. Rev. Lett. **63** (1989) 2173; B. Advera *et al.*, L3 Collab., Phys. Lett. B **231** (1989) 509; I. Decamp *et al.*, OPAL Collab., *ibid.*, **231** (1989) 519; M.Z. Akrawy *et al.*, DELPHI Collab., *ibid.*, **231** (1989) 539; C.Caso *et al.*, (Particle Data Group), Eur. Phys. J. **C 3** (1998) 1.
- [12] Z. Berezhiani and E. Nardi, Phys. Rev. D **52** (1995) 3087; C.T. Hill, E.A. Paschos, Phys. Lett. B **241** (1990) 96.
- [13] H.J. He, N. Pokorsky and S. Su, Phys. Rev. **D64** (2001) 053004; V.A. Novikov, L.B. Okun, A.V. Rozanov, and M.I. Vysotsky, Phys. Lett. **B529** (2002) 111.
- [14] Y. Nir and D. Silverman, Phys. Rev. **D42** (1990) 1477; W-S, Choong and D. Silverman, Phys. Rev. **D49** (1994) 2322; L.T. Handoko, hep-ph/9708447.
- [15] V. Barger, Y.B. Dai, K. Whisnant and B.L. Young, hep-ph/9901380; R.N. Mohapatra, hep-ph/9702229; S. Mohanty, D.P. Roy and U. Sarkar, hep-ph/9810309; S.C. Gibbons, *et al.*, Phys. Lett. **B430** (1998) 296; V. Barger, K. Whisnant and T.J. Weiler, Phys. Lett. **B427**, (1998) 97; V. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, Phys. Rev. **D58** (1998) 093016.
- [16] W.S. Hou, A. Soni and H. Steger, Phys. Lett. **B192** (1987) 441; W.S. Hou, R.S. Willey and A. Soni, Phys. Rev. Lett. **58**, 1608 (1987); Erratum, *ibid* **60**, 2337 (1987); J.F. Gunion, Douglas W. McKay, H. Pois, Phys. Lett. B **334** (1994) 339; Phys. Rev. D **51** (1995) 201.
- [17] A. Arhrib and W.S. Hou, hep-ph/0211267.
- [18] M. Aliec, A. Özpıneci and M. Savci, Nucl. Phys. **B585**, 275 (2002); hep-ph/0301078.
- [19] K. Belotsky, D. Fargion, M. Khlopov, R. Konoplich and K. Shibaev, hep-ph/0210153; K.M. Belotsky, T. Damour and M.Yu. Khlopov, Phys.Lett. **B529** (2002) 10.
- [20] LEP1.5 Collab., J. Nachtman, hep-ex/960615.
- [21] DØ Collab., S. Abachi *et al.*, Phys. Rev. Lett. **78** (1997) 3818.
- [22] C.S. Huang, W. J. Huo and Y.L. Wu, Mod. Phys. Lett. **A14** (1999) 2453.
- [23] C.S. Huang, W. J. Huo and Y.L. Wu, Phys. Rev. **D64**, 016009(2001).

- [24] W. J. Huo, Eur. Phys. J. **C24** (2002) 275.
- [25] CLEO Collaboration, M. Acciarri *et al.*, Phys. Lett. **B363**, 137(1995).
- [26] BaBar Collaboration, B. Aubert *et al.*, Phys. Rev Lett. **87**, 241803(2001).
- [27] G.-L. Lin, J. Liu, Y.-P. Yao, Phys. Rev. Lett. **64** (1990) 1498; Phys. Rev. **D42** (1990) 2314; Mod. Phys. Lett. **A6** (1991) 1333; H. Simma, D. Wyler, Nucl. Phys. **B344** (1990) 283; S. Herrlich, J. Kalinowski, Nucl. Phys. **B381** (1992) 501.
- [28] G. Hiller and E.O. Iltan, Phys. Lett. **B409** (1997) 425; C.-H. Chang, G.-L. Lin and Y.-P. Yao, **B415** (1997)395.
- [29] S.W. Bosch and G. Buchalla, *The double radiative decays $B \rightarrow \gamma\gamma$ in the heavy quark limit*, JHEP 0208 (2002) 054; S.W. Bosch, *Exclusive Radiative Decays of B Mesons in QCD Factorization*, Ph.D. Thesis, hep-ph/0208203.
- [30] M. R. Ahmady, E. Kou, A. Sugamoto, preprint hep-ph/9708347; G. Hiller, E.O. Iltan, preprint hep-ph/9708477; S. Choudlhury, J. Ellis, hep-ph/9804300.
- [31] T.M. Aliev, G. Turan, Phys. Rev. **D48** (1993) 1176; T.M. Aliev, G. Hiller, E.O. Iltan, preprint hep-ph/9708382; T.M. Aliev, E.O. Iltan, preprint hep-ph/9803459.
- [32] J.J. Cao, Z.J. Xiao and G.R. Lu, Phys. Rev. **D64** (2001)014012.
- [33] S. Bertolini, J. Matias, preprint hep-ph/9709330.
- [34] P. Singer, D.-X. Zhang, Phys. Rev. **D56** (1997) 4274.
- [35] Zhenjun Xiao, Cai-Dian Lü and Wujun Huo, hep-ph/0301221.
- [36] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. **D66**, 010001 (2002).
- [37] For a review of recent progress, see J. Flynn, in International Conference on High Energy Physics, Warsaw, Poland, 1996.

TABLES

TABLE I. The values of Wilson coefficients $C_1 - C_8$ at $\mu = m_b$ in SM

C_1	C_2	C_3	C_4	C_5	C_6	$C_7^{eff}(\text{SM})$	$C_8^{eff}(\text{SM})$
0.235	-1.100	-0.011	0.024	-0.007	0.029	0.306	0.146

FIGURES




FIG. 1. Magnetic photon and gluon penguin diagrams with the fourth generation t' quark.




FIG. 2. Branching ratios of $B_s \rightarrow \gamma\gamma$ against the mass of t' quark for $V' = V_{t's}^* V_{t'b} = -7.0 \times 10^{-2}$ and 2.5×10^{-3} respectively. The lower two horizontal lines represent the SM prediction the branching ratio.




FIG. 3. Branching ratios of $B_s \rightarrow \gamma\gamma$ as a function of $V' = V_{t's}^* V_{t'b}^{(1)}$ for $m_{t'} = 200$ GeV, 400 GeV, 600 GeV, 800 GeV and 1000 GeV, respectively. The two horizontal lines represent the SM prediction the branching ratio.




FIG. 4. Branching ratios of $B_s \rightarrow \gamma\gamma$ as a function of $V' = V_{t's}^* V_{t'b}^{(2)}$ for $m_{t'} = 200$ GeV, 400 GeV, 600 GeV, 800 GeV and 1000 GeV, respectively.




FIG. 5. Branching ratios of $B_d \rightarrow \gamma\gamma$ as a function of the mass of t' quark for $V' = V_{t'd}^* V_{t'b}$ takes several values in the range of -1.0×10^{-4} and 0.5×10^{-4} .




FIG. 6. Branching ratios of $B_s \rightarrow \gamma\gamma$ as a function of $V' = V_{t'd}^* V_{t'b}$ for $m_{t'} = 200$ GeV, 400 GeV, 600 GeV, 800 GeV and 1000 GeV, respectively.